# **CONCUSSION FACTS**

Athletes who experience one or more of the signs and symptoms listed below after a bump, blow, or jolt to the head or body may have a concussion.

## SYMPTOMS REPORTED BY ATHLETE:

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

### SIGNS OBSERVED BY COACHING STAFF:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- · Is unsure of game, score, or opponent
- · Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes
- Can't recall events prior to hit or fall
- Can't recall events after hit or fall

# IF YOU SUSPECT THAT AN ATHLETE HAS A CONCUSSION, YOU SHOULD TAKE THE FOLLOWING STEPS:

### **HEADS UP ACTION PLAN:**

- **1**. Remove the athlete from play.
- 2. Keep the athlete out of play the day of the injury.
- 3. Obtain permission from an appropriate health care professional that states the athlete can return to play.

# **IMPORTANT PHONE NUMBERS:**

EMERGENCY MEDICAL SERVICES

NAME:

PHONE: \_\_\_\_\_

### HEALTH CARE PROFESSIONAL

NAME:

PHONE:

STAFF AVAILABLE DURING PRACTICES

NAME: \_\_\_\_\_

PHONE:

STAFF AVAILABLE DURING GAMES

NAME:

PHONE: \_\_\_\_\_

JOIN THE CONVERSATION **L** www.facebook.com/CDCHeadsUp

TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).

HEADS UP CONCUSSION