

## **Concussion Protocol**

A concussion is a brain injury and all brain injuries are serious. They are caused by a bump, blow, or jolt to the head, or by a blow to another part of the body with the force transmitted to the head. They can range from mild to severe and can disrupt the way the brain normally works. Even though most concussions are mild, all concussions are potentially serious and may result in complications including prolonged brain damage and death if not recognized and managed properly. In other words, even a ding or a bump on the head can be serious. You can't see a concussion and most sports concussions occur without loss of consciousness. Signs and symptoms of concussion may show up right after the injury or can take hours or days to fully appear. If your child reports any symptoms of concussion, or if you notice the symptoms or signs of concussion yourself, seek medical attention right away.

Feeling foggy or groggy

Change in sleep patterns

Fatigue or low energy

Nervousness or anxiety

Drowsiness

Don't feel right

Amnesia

Sadness

•

•

•

•

### Symptoms may include one or more of the following:

- Headaches
- Pressure in head
- Nausea or vomiting
- Neck pain
- Balance problems or dizziness
- Blurred, double, or fuzzy vision
- Sensitivity to light or noise
- Feeling sluggish or slowed down

# Signs observed by teammates, parents and coaches include:

- Appears Dazed
- Vacant facial expression
- Confused about assignment Forgets plays
- Is unsure of game, score, or opponent
- Moves clumsily or displays incoordination
- Answers questions slowly
- Slurred speech
- Shows behavior or personality changes

### What can happen if my child keeps on playing with a concussion or returns too soon?

Athletes with the signs and symptoms of concussion must be removed from play immediately. Continuing to play with the signs and symptoms of a concussion leaves the young athlete especially vulnerable to greater injury. There is an increased risk of significant damage from a concussion for a period of time after that concussion occurs, particularly if the athlete suffers another concussion before completely recovering from the first one. This can lead to prolonged recovery, or even to severe brain swelling (second impact syndrome) with devastating and even fatal consequences. It is well known that adolescent or teenage athlete will often under report symptoms of injuries. And concussions are no different. As a result, education of administrators, coaches, parents and athletes is the key for athlete safety.

### If you think your child has suffered a concussion

Any athlete even suspected of suffering a concussion must be removed from the meet or practice *immediately.* No athlete may return to activity after an apparent head injury or concussion, regardless of how mild it seems or how quickly symptoms clear, without medical clearance. Close observation of the athlete should continue for several hours. The Zackery Lystedt Law in Washington State requires the consistent and uniform implementation of long and well established return to play concussion guidelines that have been recommended for several years: a youth athlete who is suspected of sustaining a concussion or head injury in a practice or game shall be removed from competition at that time and may not return to play until the athlete is evaluated by a licensed heath care provider trained in the evaluation and management of concussion and received written clearance to return to play from that health care provider. You should also inform your child's coach if you think that your child may have a concussion Remember it's better to miss one meet or practice than miss the whole season. And when in doubt, the athlete sits out.

For current and up-to-date information on concussions you can go to: http://www.cdc.gov/concussion/HeadsUp/youth.html

- Irritability
- More emotional
- Confusion
- Concentration or memory problems (forgetting game plays)
- Repeating the same question/comment?
- Can't recall events prior to hit
- Can't recall events after hit
- Seizures or convulsions
- Any change in typical behavior or personality
- Loses consciousness